

1.4

12. The denominator of the function f is equal to 0 when $(x - 3)(x + 1) = 0$, that is, when $x = 3$ or $x = -1$. So f is discontinuous at 3 and -1 .

24. Since $\lim_{x \rightarrow 0} (-|x| + 1) = 1$ and $f(0) = 0$, f is discontinuous at 0.

30. We require that $\lim_{x \rightarrow 2^-} (kx + 1) = \lim_{x \rightarrow 2^+} (kx^2 - 3)$, or $2k + 1 = 4k - 3$.

Solving this last equation, we obtain $k = 2$.

31. Since $\lim_{x \rightarrow 0} \frac{\sin 2x}{x} = \lim_{x \rightarrow 0} \left(2 \cdot \frac{\sin 2x}{2x}\right) = 2$, we require that $f(0) = 2$. So we take $c = 2$.

52.
$$\lim_{x \rightarrow 1} \frac{2x^3 + x - 3}{x - 1} = \lim_{x \rightarrow 1} \frac{(x - 1)(2x^2 + 2x + 3)}{x - 1} = \lim_{x \rightarrow 1} (2x^2 + 2x + 3) = 7.$$

So f will be continuous at 1 if we define $f(x) = \begin{cases} \frac{2x^3 + x - 3}{x - 1} & \text{if } x \neq 1 \\ 7 & \text{if } x = 1 \end{cases}$

54.
$$\lim_{x \rightarrow 4} \frac{4 - x}{2 - \sqrt{x}} = \lim_{x \rightarrow 4} \frac{(2 - \sqrt{x})(2 + \sqrt{x})}{2 - \sqrt{x}} = \lim_{x \rightarrow 4} (2 + \sqrt{x}) = 4.$$

So we define $f(x) = \begin{cases} \frac{4 - x}{2 - \sqrt{x}} & \text{if } x \neq 4 \\ 4 & \text{if } x = 4 \end{cases}$

60. $f(x) = x^2 - 4x + 6$ is continuous on $[0, 3]$. $f(0) = 6$ and $f(3) = 3$.

Since $f(3) \leq 3 \leq f(0)$, there exists a number c in $[0, 3]$ such that $f(c) = 3$.

To find c we solve $x^2 - 4x + 6 = 3 \Rightarrow x^2 - 4x + 3 = (x - 3)(x - 1) = 0$ giving $x = 1$ or 3 .

Therefore $c = 1$ or 3 .

64. $f(x) = x^4 - 2x^3 - 3x^2 + 7$ is continuous on $[1, 2]$.

$f(1) = 3 > 0$ and $f(2) = -5 < 0$. Therefore, by


Theorem 7, $f(x) = 0$ has at least one root in $(1, 2)$.

70. a. $\lim_{x \rightarrow 0^-} f(x) = \lim_{x \rightarrow 0^-} (-x^2) = 0$ and

$\lim_{x \rightarrow 0^+} f(x) = \lim_{x \rightarrow 0^+} (x + 1) = 1$. This shows that f is not

continuous at 0, and therefore is not continuous on $[-1, 1]$.

b. $f(-1) = -1$ and $f(1) = 2$. The number $\frac{1}{2}$ lies between $f(-1)$ and $f(1)$, but there is no c in $[-1, 1]$ such that $f(c) = \frac{1}{2}$. (See the figure.)



1.5

$$\begin{aligned} 12. \text{ a. } m_{\text{sec}} &= \frac{f(2+h) - f(2)}{(2+h) - 2} = \frac{[(2+h)^3 + (2+h)] - (2^3 + 2)}{h} \\ &= \frac{h^3 + 6h^2 + 13h}{h} = h^2 + 6h + 13 \end{aligned}$$

$$\text{b. } m_{\text{tan}} = \lim_{h \rightarrow 0} \frac{f(2+h) - f(2)}{(2+h) - 2} = \lim_{h \rightarrow 0} (h^2 + 6h + 13) = 13$$

$$\text{c. } y - 10 = 13(x - 2) \Rightarrow y = 13x - 16$$


$$\begin{aligned} 18. \lim_{h \rightarrow 0} \frac{f(4+h) - f(4)}{(4+h) - 4} &= \lim_{h \rightarrow 0} \frac{\sqrt{4+h} - \sqrt{4}}{h} = \lim_{h \rightarrow 0} \frac{(\sqrt{4+h} - 2)(\sqrt{4+h} + 2)}{h(\sqrt{4+h} + 2)} \\ &= \lim_{h \rightarrow 0} \frac{h}{h(\sqrt{4+h} + 2)} = \frac{1}{4} \end{aligned}$$

36. Using the definition of the derivative, we find $f(x) = 2x^{1/4}$ and $a = 16$.

38. Using the definition of the derivative, we find $f(x) = 2^x$ and $a = 3$.

Chapter 1 Review

4.



a. $\lim_{x \rightarrow 2^-} f(x) = -1$

b. $\lim_{x \rightarrow 2^+} f(x) = 1$

c. $\lim_{x \rightarrow 2} f(x)$ does not exist.

$$12. \lim_{x \rightarrow 3} \frac{27 - x^3}{x - 3} = \lim_{x \rightarrow 3} \frac{-(x - 3)(x^2 + 3x + 9)}{x - 3} = \lim_{x \rightarrow 3} [-(x^2 + 3x + 9)] = -27$$

$$16. \lim_{x \rightarrow 4} \frac{x - 4}{\sqrt{x} - 2} = \lim_{x \rightarrow 4} \frac{(\sqrt{x} + 2)(\sqrt{x} - 2)}{\sqrt{x} - 2} = \lim_{x \rightarrow 4} (\sqrt{x} + 2) = 4$$

$$18. \lim_{x \rightarrow 2^+} \frac{x - 2}{|x - 2|} = \lim_{x \rightarrow 2^+} \frac{x - 2}{x - 2} = 1 \text{ since } |x - 2| = x - 2 \text{ for all } x \geq 2.$$

$$\begin{aligned} 22. \lim_{x \rightarrow 0} x \cot 2x &= \lim_{x \rightarrow 0} x \cdot \frac{\cos 2x}{\sin 2x} = \lim_{x \rightarrow 0} \cos 2x \cdot \lim_{x \rightarrow 0} \frac{\frac{1}{2}(2x)}{\sin 2x} \\ &= 1 \cdot \frac{1}{2} \cdot \lim_{x \rightarrow 0} \frac{2x}{\sin 2x} = \frac{1}{2} \end{aligned}$$

42. If $x \neq 0$, then $f(x)$ is continuous. Next, observe that $-|x| \leq x \sin(1/x) \leq |x|$.

Since $|\sin(1/x)| \leq 1$, the Squeeze Theorem implies that $\lim_{x \rightarrow 0} x \sin(1/x) = 0 = f(0)$,

and so f is continuous at 0 as well. Therefore, f is continuous on $(-\infty, \infty)$.

2.1

12. $f(x) = -\frac{2}{\sqrt{x}} \Rightarrow$

$$\begin{aligned}f'(x) &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = \lim_{h \rightarrow 0} \frac{-\frac{2}{\sqrt{x+h}} - \left(-\frac{2}{\sqrt{x}}\right)}{h} \\&= 2 \lim_{h \rightarrow 0} \frac{\frac{-\sqrt{x} + \sqrt{x+h}}{\sqrt{x}\sqrt{x+h}}}{h} = 2 \lim_{h \rightarrow 0} \frac{\sqrt{x+h} - \sqrt{x}}{h\sqrt{x}\sqrt{x+h}} \cdot \frac{\sqrt{x+h} + \sqrt{x}}{\sqrt{x+h} + \sqrt{x}} \\&= 2 \lim_{h \rightarrow 0} \frac{(x+h) - x}{h\sqrt{x}\sqrt{x+h}(\sqrt{x+h} + \sqrt{x})} = 2 \lim_{h \rightarrow 0} \frac{h}{h\sqrt{x}\sqrt{x+h}(\sqrt{x+h} + \sqrt{x})} \\&= 2 \lim_{h \rightarrow 0} \frac{1}{\sqrt{x}\sqrt{x+h}(\sqrt{x+h} + \sqrt{x})} = \frac{2}{\sqrt{x}\sqrt{x}(2\sqrt{x})} = \frac{1}{x\sqrt{x}} \text{ with domain } (0, \infty).\end{aligned}$$

15. $f(x) = x^2 + 1 \Rightarrow$

$$\begin{aligned}f'(x) &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = \lim_{h \rightarrow 0} \frac{[(x+h)^2 + 1] - (x^2 + 1)}{h} \\&= \lim_{h \rightarrow 0} \frac{(x^2 + 2xh + h^2 + 1) - (x^2 + 1)}{h} \\&= \lim_{h \rightarrow 0} \frac{h(2x + h)}{h} = \lim_{h \rightarrow 0} (2x + h) = 2x\end{aligned}$$

The slope of the tangent line at $(2, 5)$ is $f'(2) = 2(2) = 4$.

An equation of the tangent line is $y - 5 = 4(x - 2)$ or $y = 4x - 3$.

$$50. \lim_{x \rightarrow 0^-} f(x) = \lim_{x \rightarrow 0^-} (x + 1) = 1, \quad \lim_{x \rightarrow 0^+} f(x) = \lim_{x \rightarrow 0^+} (x^2 + 1) = 1.$$

Therefore, $\lim_{x \rightarrow 0} f(x) = 1$. Also, $f(0) = 0 + 1 = 1$, and so $\lim_{x \rightarrow 0} f(x) = f(0)$.

Therefore, f is continuous at 0.

To show that f is not differentiable at 0, let $h < 0$ and consider

$$\lim_{h \rightarrow 0^-} \frac{f(0+h) - f(0)}{h} = \lim_{h \rightarrow 0^-} \frac{(h+1) - 1}{h} = \lim_{h \rightarrow 0^-} 1 = 1.$$

Next, if $h > 0$, then

$$\lim_{h \rightarrow 0^+} \frac{f(0+h) - f(0)}{h} = \lim_{h \rightarrow 0^+} \frac{[(0+h)^2 + 1] - 1}{h} = \lim_{h \rightarrow 0^+} h = 0.$$

This shows that $\lim_{h \rightarrow 0} \frac{f(0+h) - f(0)}{h}$ does not exist, and so by definition,

f is not differentiable at 0.

$$52. f(x) = \begin{cases} x \sin(1/x) & \text{if } x \neq 0 \\ 0 & \text{if } x = 0 \end{cases} \quad \text{Because } |\sin(1/x)| \leq 1, \text{ we have } 0 \leq |x \sin(1/x)| \leq |x|.$$

Since $\lim_{x \rightarrow 0} |x| = 0$, the Squeeze Theorem implies that $\lim_{x \rightarrow 0} x \sin(1/x) = 0 = f(0)$,

and so f is continuous at 0. To show that f is not differentiable at 0, we compute

$$f'(0) = \lim_{h \rightarrow 0} \frac{f(0+h) - f(0)}{h} = \lim_{h \rightarrow 0} \frac{h \sin(1/h)}{h} = \lim_{h \rightarrow 0} \sin(1/h)$$

which does not exist. Therefore, f is not differentiable at 0.